

SCULPTRA



PARIAESTHETICS

DISCLAIMER : THE INFORMATION PROVIDED HERE IS AT A HIGH LEVEL AND IS NOT EXHAUSTIVE. DETAILED INFORMATION WILL BE PROVIDED AT THE TIME OF YOUR CONSULTATION ALONG WITH A COMPREHENSIVE INFORMED CONSENT FORM.

PRE TREATMENT INSTRUCTIONS:

- Avoid the use of anti-inflammatory medications such as aspirin and Advil for up to 1 week pre and post injection as they will increase your risk of bruising
- Do not drink alcoholic beverages 24 hours before
- You may take Arnica tablets 2-3 days prior to your injection to reduce the risk of bruising
- Do not apply products that are potentially irritating for 2 -3 days before and after treatment (examples- tretinoin/retin-a, glycolic acid, benzoyl peroxide, hydroquinone)
- Inform your provider if you have a history of cold sores to receive advice on antiviral therapy prior to treatment
- Do not drink alcoholic beverages 24 hours before your appointment,



POST TREATMENT INSTRUCTIONS

- You can resume your activities immediately after the injection
- About 24 hours after the treatment, any swelling or redness will disappear
- Bruises fade in 2 to 3 weeks. During this period, it is recommended to avoid any sun exposure.
- You will need to massage the area of treatment for 5 minutes, 5 times a day, for 5 days. (We will show you how to massage the area after your first treatment)